



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

V. *On the length of the French Mètre estimated in parts of the English standard.* By Captain Henry Kater, F. R. S.

Read February 5, 1818.

ONE of the objects of the committee of the Royal Society appointed for the purpose of determining the length of the seconds pendulum, being the comparison of the French mètre with the British standard measure, two mètres were procured from Paris for that purpose, the one made in the usual manner and called the *mètre à bouts*, and the other a bar of platina on which the length of the mètre is shown by two very fine lines; this is named the *mètre à traits*.

The width of the *mètre à bouts* is one inch, and its thickness 0.3 of an inch. On one side the word "MÈTRE" is engraved, and on the other "FORTIN À PARIS." The terminating planes are supposed to be perfectly parallel, and the distance between them is the length of the mètre.

The *mètre à traits* is the same width as the *mètre à bouts*, but only a quarter of an inch thick. The lines expressing the length of the mètre are so fine that one of them is scarcely perceptible even with the assistance of a microscope, unless the light be very favourable. The situation of the lines may however be discovered by two strong black dots made with a graver at the extremities of each, and a fine line crosses them at right angles to indicate the parts from which the measurements are to be taken.

This mètre previous to being brought from Paris, was

compared with a standard mètre by M. ARAGO, with all that care and ability which he is so well known to possess, and which so delicate an operation requires. The result was, that the distance between the lines was found to be less than a mètre by $\frac{17.52}{1000}$ of a millimètre or ,00069 of an inch.

The same micrometer microscopes were used in the comparisons which I am about to detail, as have been already described in my account of experiments on the length of the pendulum, in the Philosophical Transactions of the present year, and as the length of the mètre is nearly 39.4 inches, I was enabled to refer it to the same divisions of Sir GEORGE SHUCKBURGH'S scale as I had employed in the measurement of the pendulum.

I commenced with the *mètre à traits*. It was placed in contact with the standard scale, their surfaces being in the same plane. An excellent thermometer was laid upon the scale, and a piece of thick leather was placed upon its bulb in order to prevent its being affected by heat from the person of the observer.

The whole was suffered to remain in this state for two or three days, after which the following observations were made at various times, the microscopes being brought alternately over the mètre and the scale. The value of each division of the micrometer is $\frac{1}{23363}$ of an inch.*

* For the manner in which this value was obtained, see page 51 of the preceding paper.

Comparison of the *mètre à traits*.

| Temperature. | Reading of the microm. at 39,4 of the scale. | Reading of the microm. at the mètre | Divisions to be deducted from 39,4 inches. | Distance in inches between the lines designating the mètre. | Correction for temperature in decimals of an inch. | Distance in inches between the lines designating the mètre, the mètre being at 32°, and the scale at 62°. |
|--|--|-------------------------------------|--|---|--|---|
| 60,0 | 85,0 | 644,5 | 559,5 | 39,37606 | ,00604 | 39,37002 |
| 60,7 | 75,5 | 639,0 | 563,5 | 39,37588 | ,00589 | 39,36999 |
| 61,7 | 69,2 | 634,0 | 564,8 | 39,37583 | ,00568 | 39,37015 |
| 62,0 | 65,0 | 630,5 | 565,5 | 39,37580 | ,00562 | 39,37018 |
| 62,4 | 61,0 | 629,0 | 568,0 | 39,37569 | ,00554 | 39,37015 |
| 62,3 | 58,7 | 629,5 | 570,8 | 39,37557 | ,00556 | 39,37001 |
| 62,2 | 58,0 | 628,0 | 570,0 | 39,37560 | ,00558 | 39,37002 |
| 62,2 | 59,0 | 625,0 | 566,0 | 39,37577 | ,00558 | 39,37019 |
| 62,1 | 59,0 | 625,5 | 566,5 | 39,37575 | ,00560 | 39,37015 |
| 58,8 | 90,0 | 638,0 | 548,0 | 39,37654 | ,00629 | 39,37025 |
| 59,0 | 83,0 | 637,0 | 554,0 | 39,37629 | ,00625 | 39,37004 |
| 59,0 | 82,0 | 636,0 | 554,0 | 39,37629 | ,00625 | 39,37004 |
| 59,2 | 83,2 | 632,0 | 548,8 | 39,37651 | ,00621 | 39,37030 |
| 59,1 | 81,0 | 632,0 | 551,0 | 39,37642 | ,00623 | 39,37019 |
| Mean | | | | | | 39,37012 |
| The distance between the lines designating the mètre was found by M. ARAGO to be too little by a quantity = ,00069 of an inch which add | | | | | | + ,00069 |
| The distance from zero to 39,4 of Sir G. SHUCKBURN'S scale is too short compared with the mean of its divisions by ,00005 of an inch which subtract* | | | | | | — ,00005 |
| Hence the length of the mètre in inches of Sir G. SHUCKBURN'S scale is | | | | | | 39,37076 |

The comparison of the *mètre à bouts*, presented considerable difficulties, which I conceive it would be of little use to detail, as the necessity of comparisons of this kind is of very rare occurrence ; I shall therefore proceed to describe the method which was at last found successful.

* See page 54.

Four rectangular pieces of brass were prepared precisely similar to those described in the account of experiments on the pendulum in the Philosophical Transactions before referred to. These were marked C, *c*, D and *d*. The perfectly plane rectangular edges of the pieces C and *c*, being placed in contact, and kept thus by means of a spring, the distance of the fine lines drawn on their surfaces, parallel and very near to the rectangular edges, was found to be 500,5 divisions of the micrometer, and the pieces D and *d* being placed in like manner in contact, the distance of the lines on their surfaces estimated in the same divisions was 456,7.

The *mètre à bouts* being placed by the side of the brass scale and in contact with it, the pieces D and *d* were applied to its extremities, the surfaces of the brass pieces being a little below the surface of the *mètre* in order to preclude any error which might have arisen from the edges of the *mètre* projecting beyond its terminating planes. Each of the brass pieces was supported in this position upon a piece of lead of a sufficient thickness, and kept in close contact with the end of the *mètre* by means of a slight spring bearing against a pin driven perpendicularly into the lead.

In order to ensure a perfect contact between each brass piece and the terminating plane of the *mètre*, a flat ruler of brass was laid upon the surface of the *mètre* so as to project beyond its extremity, and the end of the lead was elevated or depressed so that the line of light seen between the piece of brass and the ruler, the eye being level with the surface of the brass, appeared to be equal in every part, when it was inferred that the surfaces of the *mètre* and of the piece of

parts of its length for one degree of FAHRENHEIT, and as this is the expansion used by the French in adjusting the length of their *mètre*, it must be employed on the present occasion. The *mètre* being taken at 32° , the expansion for the difference between this and the temperature of measurement, must be subtracted from the apparent length of the *mètre*. The English standard temperature is 62° , therefore if the temperature of measurement be under this, the expansion of the scale for such difference of temperature must be deducted from the length of the *mètre* before obtained. These two corrections are combined in the column entitled "correction for temperature." Sir G. SHUCKBURGH's standard scale is of cast brass, and as I could not conveniently determine its actual expansion with that degree of accuracy that would have satisfied me, I have taken for it, the mean result of two experiments made on plate brass, which gave me an expansion of ,0000101 parts of its length for one degree of FAHRENHEIT. The mean of most of the experiments made on the expansion of brass gives ,0000104, and had I employed this last number instead of my own, the difference in the length of the *mètre* would have been utterly inconsiderable.

Supposing then both *mètres* to be of equal authority, we have for the length of the *mètre à traits* 39,37076, and for that of the *mètre à bouts* 39,37081 inches; the mean of which, 39,37079, may be taken for the length of the *mètre* in inches of Sir G. SHUCKBURGH's standard scale when each is brought to its proper temperature.*

* The length of the *mètre* compared with BIRD's parliamentary standard is 39,37062 inches.